

TAPHONOMY OF THE BIVALVES CONCENTRATIONS FROM CAPE MELVILLE FORMATION (EARLY MIOCENE), MOBY DICK GROUP, KING GEORGE ISLAND, WEST ANTARTICA

CASATI, R.¹; ANELLI, L.E.¹; SANTOS, P.R. dos¹

¹ Instituto de Geociências da Universidade de São Paulo, IGc/USP, SP

The Cape Melville Formation consists of about 200 m of shales and silty shales with subordinate intercalation of siltstone and fine-grained sandstone of glacial marine facies. Among the rich invertebrate fauna that includes corals, brachiopods, echinoderms and crustaceans, bivalves are an important but yet poorly studied component. We present herein the results of a qualitative and quantitative taphonomic analysis of the bivalves concentrations obtained from four geological sections from the Cape Melville Formation. 1. Penguin Rookery Section: the section consists of two assemblages represented by eight taxa (n=227) shared in two beds; the first bed (n=137, 60%) is mainly composed by remains of the shallow infaunal deposit-feeder *Ennucula* sp.1 (n=91, 66%) and subsidiary gastropods and indeterminate bivalves. *Ennucula* sp.1 is chaotically distributed (Perpendicular=37%, Oblique=31%, Concordant=30%), mainly with closed articulated valves (n=86, 95%) and few disarticulated (n=76, 84%) with good quality of preservation (n=72, 79%); the assemblage from the second bed (n=90, 40%) is mainly composed by the shallow infaunal deposit-feeder *Yoldia* sp. (n=76, 84%) and subsidiary gastropods and indeterminate bivalves. *Yoldia* sp. is preferred concordant accordingly to the bed (n=43, 57%) and most of them with closed articulated valves (n=52, 69%), almost with no fragmentation (n=64, 84%) and good quality of preservation (n=17, 22%). 2. Hard Ground Section: the section consists of an assemblage represented by seven taxa (n=211) where stands out the shallow infaunal deposit-feeder *Ennucula* sp.2 (n=130, 62%) and subsidiary gastropods, corals and indeterminate bivalves. *Ennucula* sp.2 is chaotically distributed (Perp.=30%, Obl.=36%; Conc=39%), preferred closed articulated (n=117, 90%), almost with no fragmentation (n=94, 72%) and with good quality of preservation (n=59, 45%). 3. Chamine Section: the section consists of an assemblage represented by seven taxa (n=69) where stands out the infaunal deposit-feeder *Neilo* (*Neilo*) sp. (n=22, 32%), the deep infaunal suspension-feeder *Panopea* (*Panopea*) cf. *Panopea* sp. (n=10, 14%) and subsidiary gastropods, crustaceans and indeterminate bivalves. *Neilo* (*Neilo*) sp. is preferred concordant (n=11, 50%), closed articulated (n=16, 73%), with no fragmentation and good quality of preservation; *Panopea* (*Panopea*) cf. *Panopea* sp. is preferred perpendicular (n=8, 80%), with closed articulated valves (n=9, 90%), no fragmentation and good quality of preservation. 4. Lava Crag Section: the section consists of an assemblage represented by four taxa (n=40) where are included infaunal deposit-feeder *Neilo* (*Neilo*) sp. (n=17, 43%) and subsidiary crabs and indeterminate bivalves. *Neilo* sp. is preferred concordant (n=12, 71%) and all the specimens are disarticulated, with no fragmentation and good quality of preservation. Therefore, based on the taphonomic characterization above, the main objective of this study is to elucidate if these accumulations were controlled by paleoecologic, biostratigraphic and/or diagenetic patterns. The Penguin Rookery and Hard Ground Sections are very close for its faunal composition and taphonomic characteristics, as well as Chamine and Lava Crag Sections. This sort of study is essential for the comprehension of the genesis of the fossil concentrations and for modern paleobiology analysis.